

Depression, Psychosis & Suicide Ideation - The Dark side of Pfizer & Moderna COVID-19 mRNA Vaccine effects on the BRAIN. I review 9 studies on Suicidality, 12 on Psychosis, 5 on Depression!



DR. WILLIAM MAKIS MD

AUG 19, 2023 • PAID



42



6



3

Share



COVID-19 mRNA vaccine spike protein accumulates in the brain and causes many types of neurological and psychiatric injuries. These are

being suppressed by the medical establishment and the media.

Here is the latest published literature:

Suicidal Ideation and Suicide Attempts after mRNA:

One of my most controversial hypotheses is that COVID-19 mRNA vaccine spike protein accumulates in the brain and causes a severe inflammatory and immune reaction which leads to neurological and psychiatric adverse events, and increased risk of suicidal ideation.

However, there is growing literature to support this:

Study 1 - CDC confirms in Aug.2021: Two teenagers committed suicide after Pfizer COVID-19 mRNA vaccine

- ([Hause et al](#)) - CDC reviewed reports of death after Pfizer mRNA
- Out of 14 teenager deaths reviewed by CDC, two were suicides
- ([Hesse et al](#)) - confirmed the findings

Study 2 - 12% of Pakistani Dental College staff had suicidal thoughts for several days after COVID-19 vaccination, 1.6% had suicidal thoughts almost every day!

- ([Maruf et al](#)) - Pakistani questionnaire study on 250 Dental College staff after they were COVID-19 vaccinated, to study mental health impact:
- 11.6% (29/250) had suicidal thoughts for a few days
- 1.6% (4/250) had suicidal thoughts nearly every day!

Study 3 - 45 yo Latvian man had psychosis & suicide attempt after mRNA

- ([Renemane et al](#)) - 45 yo Latvian man had 2nd dose of mRNA vaccine
- immediately developed insomnia, anxiety, tremors
- 2 weeks after jab “discovered white powder under the carpet in his apartment”

- 1 month after jab he was “receiving commands” including to commit suicide
- he attached a rope, tried to hang himself but his father stopped him.

Study 4 - 40 yo white man became suicidal after Moderna

- ([Guina et al](#)) - 40 yo white man had 2nd Moderna dose
- post jab: he had depressed mood, suicidal ideation and auditory hallucinations encouraging him to kill himself
- he lost his job after a verbal altercation with his boss, prompting him to seek hospitalization
- on admission he was hyperactive, impulsive, irritable, flight of ideas, pressured and hyperverbal speech, poor memory

Study 5: 37 yo Japanese man had psychosis & suicide attempt after Moderna mRNA

- ([Kita et al](#)) - 37 yo Japanese man with no psychiatric history had Moderna booster
- He presented 4 days later with talkativeness and grandiose delusions, saying that he had won 2 billion yen in horse racing. Had emotional instability (crying), sleeplessness, excitement, hyperactivity and sexual deviance.
- He was discharged. Nine days after Moderna booster he jumped from the 2nd floor of his house, and was brought back to hospital by ambulance.
- He exhibited flight of ideas, hyperactivity, distraction, hyperthymia and religious delusions, such as saying “my child is God”. He displayed lack of insight and became enraged when his actions were restricted.
- Diagnosis: acute mania with psychotic features, stayed 66 days in the hospital.

Study 6: 57 yo Turkish man had suicide attempt after 2nd Pfizer dose

- ([Yesilkaya et al](#)) - 57 year old man presented to ER with irritability, sleeplessness, talking to himself and suicidal attempt with thoughts of extinction 3 days after receiving 2nd dose of Pfizer.

- had increased speech output, speed and psychomotor activity
- had no history of psychiatric illness

Study 7: 45 yo man had psychosis & suicide attempt after 2 Pfizer doses

- ([Deng et al](#)) - 45 yo man had 2 Pfizer mRNA doses in May 2021.
- In June 2021 he presented to ER with depression, paranoia, intermittent agitation and had preliminary diagnoses of bipolar disorder and delirium
- 2 months after his 2 Pfizer doses he attempted to commit suicide by drowning himself in a toilet. He was feeling depressed and suicidal
- also experienced auditory hallucinations, paranoia and persecutory delusions.

Study 8: 26 year old woman had 1st Pfizer dose, developed a severe panic disorder & suicidal ideation

- ([de Souza Melo](#)) - 26 year old woman had 1st Pfizer dose
- she was admitted to psychiatry with suicidal ideation after episodes of severe and recurrent panic attacks
- she fully recovered, then had 2nd Pfizer dose, and her panic attacks returned.

SUICIDE IDEATION & ATTEMPTS BY KIDS IN VAERS:

VAERS 1406268: 13 yo boy (Nebraska)

One day after 1st dose of Pfizer he developed headaches, extreme fatigue, disconnect with reality. He stopped communicating, developed paranoia, began to have self-neglect, not caring for self or doing normal hygiene. Within days had visual and tactile hallucinations, flat affect. Began cutting himself.

VAERS 1513678: 12 yo girl (New Mexico)

She reported having **suicidal thoughts, hearing voices**. Patient started **self-harming** after 1st dose of Pfizer. After 2nd dose of Pfizer, the self-harm got worse and she became very depressed.

VAERS 1980939: 11 yo boy (Florida)

On same day after 2nd dose of Pfizer, he developed severe headache, fever, sleepwalked. At one point he **believed he was possessed, was hallucinating and wanted to jump out the window**.

VAERS 2043532: 12 yo boy (Washington)

Got 3rd dose of Pfizer, next day he had 5 min episodes of hallucinations in which he says and does things that do not make sense. At 1am woke up, extremely agitated, yelling, screaming, crying, heart racing, **started saying he was going to die or needed to die**. He is not lucid.

When asked questions he does not make sense or does not know basic information such as his age or birthdate. **He walked to the banister from his room on the 2nd story and began climbing over the edge. My husband grabbed him and took him to his bed** holding him down and asking him to count to 10. He slowly begins to calm and all of a sudden becomes lucid again and starts making sense. He has no memory if his actions during the hallucinations.

He began having a 2nd hallucination. He enters he hallways and began yelling at his brother as if he was there, but his brother was in his room sleeping. He ran downstairs to my bedroom and **started yelling he was going to die. I walked him back upstairs and he told my husband to kill him. When we asked him what he did that day he said he jumped over the edge and jumped off a cliff.**

=====

PSYCHOSIS

Study 1 - 60 yo African American woman post 2nd Pfizer mRNA dose had Bipolar Disorder exacerbation (psychosis)

- ([Guina et al](#)) - 60 yo woman had 2nd Pfizer dose, within 1 week she had anxiety, insomnia
- on admission: increased energy, impulsivity, verbose speech, flight of ideas, paranoia, ideas of reference - diagnosed with psychotic episode

Study 2 - 20 yo woman had psychosis after Pfizer mRNA

- ([Flannery et al](#)) - 20 yo presented to ER 1 week after 1st Pfizer dose
- had auditory hallucinations
- one morning she removed her clothes and had a bowel movement on the floor, then became increasingly psychotic

Study 3 - 38 yo Brazilian woman had refractory psychosis after Pfizer mRNA

- ([Neves et al](#)) - had 1st Pfizer dose, presented to ER within 24hr
- She became delusional, aggressive, disheveled with poor hygiene, unfriendly face, lacking self-orientation and uncooperative with doctors & staff.
- She believed staff were persecuting her, that several city mayors were her parents. Her mood was dysphoric and enraged.
- She was put on psychiatric medications but didn't improve.
- After 6 months of hospitalization, she remained hostile, persecutory and refusing to talk to staff and her psychosis was defined as refractory. She never recovered.

Study 4: 15 yo boy had psychosis after Pfizer mRNA

- ([Yen-Li Lien et al.](#)) 15 yo Taiwanese boy presented to ER with agitation, involuntary limb stretching and screaming 2 days after 2nd dose of Pfizer
- After admission, he had bizarre behaviors, including sitting up and lying down frequently and taking up the mannerism of praying in bed.
- had auditory hallucinations and delusions, persisted > 1 month after discharge

Study 5: 31 yo man had psychosis after Moderna mRNA

- ([Reinfeld et al.](#)) A 31-year-old, single Hispanic man had two doses Moderna
- brought to ER by police
- was anxious, guarded, superficial and grandiose.
- He reported becoming 'clairvoyant', being able to talk with dead people, hearing 'people drumming outside his house' and the constant voice of a co-worker he believed was his lover

Study 6: 45 yo Asian-American married woman, had psychosis after Moderna mRNA

- ([Alphonso et al.](#)) 45 yo woman with no psychiatric history, presented to psychiatry after having psychosis for 3 months.
- Her new-onset paranoia and auditory hallucinations began 1 month after receiving her second Moderna COVID-19 vaccine. She abruptly quit her job of 18 years and stopped eating—losing 56 lb in 4 months.
- The patient would only get dressed in the dark, believing people could see her in her closet or bathroom behind a closed door. She would pace all day and stopped doing the household chores she had done for years.
- She would wake her husband in the middle of the night trying to convince him that the neighbors were going to hurt her and were breaking into their home.
- Due to this irrational fear, she would sleep only 2 to 3 hours per night and no longer slept in the bedroom, preferring to lock herself in the bathroom to sleep.
- She heard whispers of people talking about her and would carry on full conversations with them. She would only do this in a room alone with the door closed.

Study 7: 42 yo Turkish man had psychosis after Pfizer

- ([Yesilkaya](#)) - 42 yo man with no psychiatric history was admitted to psychiatric ER 5 days after 1st dose of Pfizer with complaints of irritability and sleeplessness.
- The patient had delusions that his family was being followed by the deep state and that they were in danger.

- Physical exam: **anxious, dysphoric mood, persecutory and reference delusions**
- Diagnosed with acute mania / psychotic episode

Study 8: 26 yo woman has acute psychosis & lupus after Pfizer mRNA

- ([Fekih-Romdhane et al](#)) - 26 yo woman had 2nd Pfizer dose
- 4 days later presented to ER with **acute psychomotor agitation, incoherent speech and total insomnia evolving for 5 days**
- **she was diagnosed with a brief psychotic disorder**
- was also found to have an acute SLE (lupus) cerebritis

Study 9: 20 year old woman in Saudi Arabia had psychosis after COVID-19 mRNA vaccine

- ([Aljeshi et al](#)) - 20 year old College student had 2nd mRNA vaccine (Saudi Arabia gives Pfizer & Moderna mRNA vaccines).
- she presented to ER with 4 weeks of anxiety, **sleep disturbance, restlessness, reduced appetite that started 4 days after her 2nd mRNA jab.**
- in ER, she complained of “**hearing terrifying voices and seeing random people stare at her**”
- in hospital she became **agitated, aggressive, disoriented and complained of auditory and visual hallucinations.** She had disjointed speech and grossly disorganized and disruptive behavior and was disoriented
- she was diagnosed with **acute psychosis** and stayed in hospital 28 days

Study 10: 41 yo man, 35 yo man develop psychosis after Pfizer mRNA

- ([Borovina et al](#)) - 41 yo male had 2nd dose of Pfizer mRNA
- 5 days later, had headaches, **paranoid delusions and severe anxiety**
- in ER he had a **depressed, irritable mood and olfactory hallucinations**

- he was discharged with **ongoing psychotic symptoms** and had to be re-admitted with changes to his anti-psychotic medications
- A 35 yo male had 1st dose of Pfizer, 3 days later he had headaches
- 2 weeks later presented to ER with **persecutory delusions and delusions of reference**
- in ER he was **agitated, dysphoric and paranoid**.
- he was discharged with partial remission of psychotic symptoms

Study 11: 32 yo man had psychosis after Moderna mRNA

- ([Al-Mashdali et al](#)) - 32 yo man had 1st dose of COVID-19 Moderna mRNA
- within 24hr he developed acute confusion, memory disturbances and auditory hallucinations
- patient improved with steroids

Study 12: Review of UK adverse events to Pfizer mRNA finds 176 psychiatric adverse events

- ([Peter-Lloret et al](#)) conducted a review of adverse psychiatric reactions to Pfizer mRNA vaccines in the UK
- Found 176 psychiatric AEs including anxiety (#1), hallucinations (#2), poor quality sleep (#3), delirium

DEPRESSION & SLEEP IMPAIRMENT:

Study 1 - Israeli study finds 11% have clinical depression after COVID-19 vaccination

- ([Greenblatt-Kimron et al](#)) - A questionnaire of 938 older Israeli adults was done
- After COVID-19 vaccination, 11% were found to have clinical depression
- ([Hoffman et al](#)) - these results were published again

Study 2 - North Carolina survey of 1245 healthcare workers

- (Kadali et al) - A survey of 1245 North Carolina healthcare workers was done for side effects of Pfizer mRNA vaccine
- 12% reported psychiatric side effects: 5% decreased sleep, 2.5% anxiety, 2% increase in sleep, 1% decrease in memory, 1% psychological stress, 0.4% depression, 0.4% manic mood

Study 3 - 38% of Pakistani Dental College staff had trouble sleeping after COVID-19 vaccination

- ([Maruf et al](#)) - Pakistani questionnaire study on 250 Dental College staff after they were COVID-19 vaccinated, to study mental health impact:
- 4.4% had trouble sleeping nearly every day
- 7.6% had trouble sleeping more than half of all days
- 38% had trouble sleeping for a few days

Study 4 - 48 year old Indian man had panic disorder & sleep impairment after Pfizer mRNA

- ([Uvais](#)) - 48 year old Indian man developed new onset panic disorder following 2nd dose of Pfizer.
- quit his job in the middle east and returned to India
- presented to ER with palpitations, fear of going crazy, severe sleep impairment

Study 5 - 59 year old woman had 4 months insomnia after Moderna mRNA

- ([Aryal et al](#)) - 59 year old woman developed 4 months of insomnia after two doses of Moderna COVID-19 mRNA vaccine
- insomnia impaired her daily life activities to the point where she sought hospitalization for it

=====

THEORIES ON NEUROLOGICAL & PSYCHIATRIC INJURY:

- ([Yesiklaya et al](#)): “One hypothesis is that SARS-CoV-2 (spike protein) can damage the central nervous system via autoimmune mechanisms due to excessive production and release of pro-inflammatory chemokines and cytokines, particularly TNF- α , IL-1 and IL-6. mRNA vaccines contain nucleotides from the genetic code of the virus that encode a viral protein. This (spike) protein is a viral antigen that can cause neuropsychiatric symptoms such as autoimmune psychosis (anti-NMDAR, AMPAR, CAPR2 encephalitis) by rapidly increasing a pro-inflammatory response and activated autoimmune mechanism ([Pollak et al., 2020](#); [Sen et al. 2021](#)). Another hypothesis suggests that the inflammatory mechanisms caused by COVID-19 can trigger neuropsychiatric symptoms via thiamine deficiency. Therefore, vaccination may also lead to neuropsychiatric symptoms by damaging thiamine metabolism ([Branco de Oliveira et al., 2021](#); [de Oliveira et al., 2021](#)).
- ([Abdelhady et al](#)): “Dutta et al. reported 19,529 neurological adverse events after COVID-19 vaccination, including encephalitis. The diagnosis of mRNA-1273 vaccine-induced encephalitis was made by Fan et al. in several recent cases. The underlying mechanism of such symptomatology is not clearly understood; some researchers theorized that SARS-CoV-2 spike protein produced by mRNA-based vaccines may act as a catalyst for the inflammatory processes that ensue, particularly in autoimmune encephalitis”
- ([Nagamine](#)): “Although the mRNA vaccine promotes the synthesis of viral spike proteins, they do not activate ACE2 receptors, and thus vaccine-induced neuroleptic malignant syndrome is a different mechanism than neuroleptic malignant syndrome caused by COVID-19 infection. After vaccination, immune-transducing cells called dendritic cells begin to work, and synthesizing proinflammatory cytokines [6]. Elevated levels of proinflammatory interleukins such as IL-6 and TNF- α have been reported in neuroleptic malignant syndrome [7]. Thus, the immunostimulatory effect of the vaccine may cause the pathogenesis of neuroleptic malignant syndrome.”
- ([Zlotnik et al](#)): “The pathophysiology of those complications is still not well understood and only based on hypotheses. In general, vaccinations can cause a strong expression of proinflammatory cytokines and a T cell response (16, 19, 20).

After vaccination, antigens are recognized as potential pathogens by both conserved pathogen- and damage-associated molecular patterns as well as pattern-recognition receptors that are found on local or peripheral circulating immune cells (eg monocytes and macrophages) and on resident stromal cells (20). **Transcription and induction of many target genes occurs, resulting in synthesis and release of pyrogenic cytokines (ie interleukin [IL]-1, IL-6, tumor necrosis factor-alpha [TNF- α], and prostaglandin-E2) into the bloodstream.** These may mimic the response to natural infection. **Following stimulation, the immune system initiates a complex series of innate immune events, releasing mediators and products of inflammation to the circulation that can affect other body systems to cause systemic side-effects, and ultimately can cause neuroinflammation in some subjects after microglia activation,** depending on the immunogenetic background and the innate immune memory (19, 20).

Regarding mRNA vaccines, it has been proposed that **spike protein expression by human cells (after translation of the mRNA COVID-19 vaccine by human cells) might trigger an inflammatory reaction, similar to that induced by the virus itself, leading to those neurological complications (17, 21).** However, other mechanisms might be involved.

My Take...

Mental Health injuries and suicides are skyrocketing in the COVID-19 mRNA vaccinated and NO ONE is willing to talk about it.

At least 11 fully COVID-19 mRNA vaccinated Canadian doctors have committed suicide, and it's being covered-up by the Canadian Medical Association, who deleted the dead doctors' "In Memoriam" pages from their own website.

CDC confirms two teenagers died by suicide after taking Pfizer. There are many published reports on suicides after Pfizer and Moderna mRNA vaccines. Doctors are not warning their patients about this risk!

Pakistani Dental College staff report after COVID-19 vaccination, 12% have some suicidal thoughts, 1.6% have suicidal thoughts on an almost daily basis!

There are published reports on **psychosis after Pfizer and Moderna mRNA vaccines**. These are often associated with **visual or auditory hallucinations, delusions, paranoia, agitation, irritability, headaches and sleep disturbances**.

Those who develop a headache after taking Pfizer or Moderna are at higher risk of a psychiatric injury or complication.

An Israeli study found 11% of older people have clinical depression after COVID-19 vaccination.

A survey of North Carolina healthcare workers found 12% had psychiatric adverse events after COVID-19 vaccination.

About 4-5% of COVID-19 vaccinated have sleep disturbances on a regular basis.

There are many scientists around the world who have published their hypotheses about how the COVID-19 vaccine spike protein causes dangerous inflammation in the brain through the production of cytokines and chemokines (signaling molecules), stimulation of the immune system, and complex autoimmune processes.



Recommend COVID Intel - by Dr. William Makis to the readers of Sunfellow on Substack

In depth intel on COVID-19, sudden deaths, mRNA vaccines, vaccine injuries, new pandemics and more...

Recommend



42 Likes · 3 Restacks

6 Comments



Write a comment...



Kamikaze ~ Spirit Wind 6 hrs ago

Just saw this yesterday. Actor Chris Peluso dies at age 40. If it could be suicide:

'A year ago, it was publicly announced that the prolific actor had stepped away from performing to seek treatment for SCHIZOAFFECTIVE DISORDER. Peluso's family confirmed his death to Playbill this week but did not disclose his cause of death.

<https://www.latimes.com/entertainment-arts/story/2023-08-18/chris-peluso-death-broadway-actor>

♡ LIKE (6) 💬 REPLY ...



Anne 5 hrs ago

This is truly frightening. I remain thankful that somehow, by the grace of God, my husband and I resisted the pressure and did not take these shots. I avoid medication as much as possible (take no prescription meds at age 71), as I had a terrifying hallucination after taking a med for one day (Amantadine, prescribed for symptoms similar to MS, which have since resolved). I've said many times that I might have killed myself, had I continued taking it. In addition to the types of cases listed in this post, Brianne Dressen, vaccine injured in a clinical trial, personally knows of people whose injuries were so bad they couldn't tolerate it, and committed suicide. She said in an interview that she considered it for awhile, but decided she could not do that to her husband and children.

♡ LIKE (4) 💬 REPLY ...

4 more comments...

